WO 2005/084680 PCT/US2005/006216

## **AMENDED CLAIMS**

received by the International Bureau on 08 August 2005 (08.08.2005): original claims 1-7 have been replaced by amended claims 1-7.

## What is claimed is:

- 1. A composition comprising a reduced isoalpha acid (RIAA) and isoalpha acid (IAA) isolated from hops, wherein the RIAA and IAA are in a ratio of about 3:1 to about 1:10 and wherein said RIAA and IAA individually comprise at least 0.1% of the composition.
- 2. The composition of claim 1, wherein said isoalpha acid is selected from isohumulone, isocohumulone, and isoadhumulone.
- 3. The composition of claim 1, wherein said reduced isoalpha acid is selected from dihydro-isohumulone, dihydro-isocohumulone, and dihydro-adhumulone.
- 4. A method for reducing PGE2 mediated inflammation, comprising administering a composition comprising a reduced isoalpha acid (RIAA) and isoalpha acid (IAA) isolated from hops, wherein the RIAA and IAA are in a ratio of about 3:1 to about 1:10 and wherein said RIAA and IAA individually comprise at least 0.1% of the composition.
- 5. The method of claim 4, wherein said isoalpha acid is selected from isohumulone, isocohumulone, and isoadhumulone.
- 6. The method of claim 4, wherein said reduced isoalpha acid is selected from dihydro-isohumulone, dihydro-isocohumulone, and dihydro-adhumulone.

WO 2005/084680 PCT/US2005/006216

7. A method FOR reducing PGE2 mediated inflammation, comprising administering at least two compounds of Genus A having the formula:

wherein R' is selected from the group consisting of carbonyl, hydroxyl, OR, and OCOR, wherein R is alkyl;

and wherein R" is selected from the group consisting of CH(CH<sub>3</sub>)<sub>2</sub>, CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, and CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, wherein the two compounds are in a ratio of about 10:1 to about 1:10 and wherein said RIAA and IAA individually comprise at least 0.1% of the composition.